ELSEVIER

Fluid Phase Equilibria 119 (1996) 241–242

FLUID PHASE EQUILIBRIA

Contents of Volume 119 (1996)

Vol. 119, Nos. 1–2

Theoretical papers

The isochoric-, isobaric- and saturation-heat capacities of the Lennard–Jones fluid from equations of state and Monte Carlo simulations
D. Boda, T. Lukács, J. Liszi and I. Szalai (Veszprém, Hungary) 1–16

Fluid phase behavior of water — non polar mixtures. Concentration Fluctuations in H_2O–X ($X = H_2$, He, Ar)
O.H. Scalise (La Plata, Argentina) and M. Silbert (Norwich, UK) 17–25

Identification of variant and invariant properties in the thermodynamics of mixtures: tests for models and computer codes
M.S. Zabaloy and J.H. Vera (Montreal, Canada) 27–49

Equations of state for associating fluids (based on the analytic chain association theory)
H. Touba and G.A. Mansoori (Chicago, IL, USA) 51–65

A hybrid equation of state for Stockmayer pure fluids and mixtures
C. Kriebel, A. Müller, J. Winkelmann (Merseburg, Germany) and J. Fischer (Wien, Austria) 67–80

Thermodynamics of branched alcohols. I. Extension of DISQUAC to *tert*-alcohols–*n*-alkanes or *tert*-alcohols–cyclohexane mixtures
J.A. González, I. García de la Fuente, J.C. Cobos (Valladolid, Spain) and U. Domańska (Warsaw, Poland) 81–96

Deviations from corresponding-states behavior in the vapor–liquid equilibrium of the square-well fluid
A. Gil-Villegas, F. Del Río (México, México) and A.L. Benavides (León, México) 97–112

Experimental papers

Vapor–liquid equilibria, activity coefficients at infinite dilution and heats of mixing for mixtures of N-methyl pyrrolidone-2 with C5 or C6 hydrocarbons and for hydrocarbon mixtures
K. Fischer and J. Gmehling (Oldenburg, Germany) 113–130

Solid–liquid equilibria for systems containing long-chain 1-alkanols. I. Experimental data for 1-dodecanol, 1-tetradecanol, 1-hexadecanol, 1-octadecanol or 1-icosanol–benzene or –toluene mixtures. Characterization in terms of DISQUAC
U. Domańska (Warsaw, Poland) and J.A. González (Valladolid, Spain) 131–151

Liquid–liquid equilibria for the system naphtha reformate–dimethyl sulphoxide
T.A. Al-Sahhaf and E. Kapetanovic (Safat, Kuwait) 153–163

Solubility of gases in butanols. II. Solubilities of nonpolar gases in 2-methyl–1-propanol from 263.15 to 303.15 K at 101.33 kPa partial pressure of gas
J. Pardo, M.C. López, J. Santafé, F.M. Royo and J.S. Urieta (Zaragoza, Spain) 165–173

Excess molar enthalpies of mixing of 1-propanol or 2-propanol with aromatic hydrocarbons at 308.15 K in terms of an association model
K.C. Singh, K.C. Kalra, S. Maken and V. Gupta (Rohtak, India) 175–190

Liquid–liquid–solid equilibria for the ternary systems water–lithium chloride–1-propanol or 2-propanol at 25 °C
V. Gomis, F. Ruiz, J.C. Asensi and P. Cayuela (Alicante, Spain) 191–195

Isobaric heat capacity of natural gases. Measurements and modelling
A. Barreau, P. Janneteau and K. Gaillard (Rueil-Malmaison, France) 197–212

In situ determination of high pressure phase diagrams of methane–heavy hydrocarbon mixtures using an infrared absorption method
Ph. Marteau, P. Tobaly (Villetaneuse, France), V. Ruffier-Meray and A. Barreau (Rueil-Malmaison, France) 213–230

Corrigendum 231

Corrigendum 233

Erratum 235

Subject index 237

Author index 239

Contents of volume 241–242